

EPA-452/R-12-003
June 2012

Regulatory Impact Analysis for the Proposed Revisions to the National Ambient Air Quality
Standards for Particulate Matter

Contract No. EP-W-11-029
Work Assignment Number: 0-03

U.S. Environmental Protection Agency
Office of Air Quality Planning and Standards
Health and Environmental Impacts Division
Research Triangle Park, NC 27711

Table 5-6. Health Endpoints and Epidemiological Studies Used to Quantify Health Impacts in the Main Analysis^a

Endpoint	Study	Study Population	Risk Estimate (95th Percentile Confidence Interval) ^a
Premature Mortality			
Premature mortality—cohort study, all-cause	Krewski et al. (2009)	>29 years	RR = 1.06 (1.04 – 1.06) per 10 µg/m ³
	Laden et al. (2006)	>24 years	RR = 1.16 (1.07 – 1.26) per 10 µg/m ³
Premature mortality, total exposures	PM _{2.5} Expert Elicitation (Roman et al., 2008)	>24 years	Varies by expert
Premature mortality— all-cause	Woodruff et al. (1997)	Infant (<1 year)	OR = 1.04 (1.02 – 1.07) per 10 µg/m ³
Chronic Illness			
Nonfatal heart attacks	Peters et al. (2001)	Adults (>18 years)	OR = 1.62 (1.13 – 2.34) per 20 µg/m ³
	<i>Pooled estimate:</i>		
	Pope et al. (2006)		β = 0.00481 (0.00199)
	Sullivan et al. (2005)		β = 0.00198 (0.00224)
	Zanobetti et al. (2009)		β = 0.00225 (0.000591)
Zanobetti and Schwartz (2006)		β = 0.0053 (0.00221)	
Hospital Admissions			
Respiratory	Zanobetti et al. (2009)—ICD 460-519 (All respiratory)	>64 years	β=0.00207 (0.00446)
	Moolgavkar (2000)—ICD 490–496 (Chronic lung disease)	18–64 years	1.02 (1.01–1.03) per 36 µg/m ³
Cardiovascular	Babin et al. (2007)—ICD 493 (asthma)	<19	β=0.002 (0.004337)
	<i>Pooled estimate:</i>		
	Zanobetti et al. (2009)—ICD 390-459 (all cardiovascular)	>64 years	β=0.00189 (0.000283)
	Peng et al. (2009)—ICD 426-427; 428; 430-438; 410-414; 429; 440-449 (Cardio-, cerebro- and peripheral vascular disease)		β=0.00068 (0.000214)
	Peng et al. (2008)—ICD 426-427; 428; 430-438; 410-414; 429; 440-449 (Cardio-, cerebro- and peripheral vascular disease)		β=0.00071 (0.00013)
	Bell et al. (2008)—ICD 426-427; 428; 430-438; 410-414; 429; 440-449 (Cardio-, cerebro- and peripheral vascular disease)		β=0.0008 (0.000107)
Asthma-related ER visits	Moolgavkar (2000)—ICD 390–429 (all cardiovascular)	20–64 years	RR=1.04 (t statistic: 4.1) per 10 µg/m ³
	<i>Pooled estimate:</i>		
	Mar et al. (2010)	All ages	RR = 1.04 (1.01 – 1.07) per 7 µg/m ³
	Slaughter et al. (2005)		RR = 1.03 (0.98 – 1.09) per 10 µg/m ³

(continued)

Table 5.B-7. Unit Values for Hospital Admissions in BenMAP 4.0.51 (Abt Associates, 2011)^a

End Point	ICD Codes	Age Range		Mean Hospital Charge (2000\$)	Mean Length of Stay (days)	Total Cost of Illness (Unit value in 2000\$)
		min.	max.			
HA, All Cardiovascular	390–429	18	64	\$26,654	4.12	\$27,119
HA, All Cardiovascular	390–429	65	99	\$24,893	4.88	\$25,444
HA, All Respiratory	460–519	65	99	\$20,667	6.07	\$21,351
HA, Asthma	493	0	64	\$9,723	3.00	\$10,051
HA, Chronic Lung Disease	490–496	18	64	\$12,836	3.90	\$13,276

^a National average median daily wage is \$112.86 (2000\$).

Table 5.B-8. Unit Values for Hospital Admissions in BenMAP 4.0.43 (Abt Associates, 2010)^a

End Point	ICD Codes	Age Range		Mean Hospital Charge (2000\$)	Mean Length of Stay (days)	Total Cost of Illness (Unit value in 2000\$)
		min.	max.			
HA, All Cardiovascular	390–429	20	64	\$22,300	4.15	\$22,778
HA, All Cardiovascular	390–429	65	99	\$20,607	5.07	\$21,191
HA, All Respiratory	460–519	65	99	\$17,600	6.88	\$18,393
HA, Asthma	493	0	64	\$7,448	2.95	\$7,788
HA, Chronic Lung Disease	490–496	20	64	\$10,194	\$5.92	\$15,375

^a National average median daily wage is \$115.20 (2000\$).

Table 5.B-9. Change in Monetized Hospitalization Benefits for 12/35

Endpoint	2000 AHRQ (millions of 2006\$)	2007 AHRQ (millions of 2006\$)	Percent Change
Respiratory hospital admissions	\$2.3	\$2.4	3.4%
Cardiovascular hospital admissions	\$3.1	\$3.2	2.1%
Work loss days	\$6.7	\$6.7	0.02%

* All estimates rounded to two significant digits.

5.B.5 Long-term PM_{2.5} Mortality Estimates using Cohort Studies in California

In Chapter 5, we described the multi-state cohort studies we used to estimate the PM_{2.5}-related mortality (i.e., Krewski et al., 2009; Laden et al., 2006), as well as summarized the effect estimates for additional cohort studies. In this appendix, we provide additional information

about cohort studies in California.¹ As shown in Table 5.x in the health benefits chapter, a large percentage of the monetized human health benefits associated with the illustrative control strategy to attain the alternative combination of standards are projected to occur in California. Specifically, for an annual PM_{2.5} standard of 12 µg/m³ in conjunction with retaining the 24-hour standard of 35 µg/m³, 70% of the total monetized benefits were estimated to occur in California and 98% for an annual PM_{2.5} standard of 13 µg/m³. For this reason, we determined that it was appropriate to consider the sensitivity of the benefits results using effect estimates for cohorts in California specifically. Although we have not calculated the benefits results using these cohort studies, it is possible to use the effect estimates themselves to determine how much the monetized benefits in California would have changed if we used effect estimates from the California cohorts. Each of the California cohort studies are summarized in the PM ISA (and thus not summarized here) with the exception of the Ostro et al. (2010, 2011) studies, which we describe below. Table 5.B.10 provides the effect estimates from each of these cohort studies for all-cause, cardiovascular, cardiopulmonary, and ischemic heart disease (IHD) mortality for each of the California cohort studies.

Ostro et al. (2010) characterize the risk of premature death associated with long-term exposure to PM_{2.5} in California among a cohort of about 134,000 current and former female public school professionals (i.e., the California Teacher's Study (CTS)). In this prospective cohort study, Ostro and colleagues estimated long-term PM exposure to several PM constituents, including elemental carbon, organic carbon, sulfates, nitrates, iron, potassium, silicon and zinc. In an erratum, Ostro et al. (2011) modified their approach to assigning PM_{2.5} levels to the cohort populations, noting that they "reanalyzed the CTS data using time-dependent pollution metrics—in which the exposure estimates for everyone remaining alive in the risk set were recalculated at the time of each death—in order to compare their average exposures up to that time with that of the individual who had died. In this way, decedents and survivors comprising the risk set had similar periods of pollution exposure, without subsequent pollution trends influencing the surviving women's exposure estimates." This change in assumption attenuated the hazard ratios significantly, though hazard ratios remained significant for cardiovascular mortality and total PM_{2.5} mass and certain constituents, nitrate and sulfate; no association was observed between all-cause mortality and total PM_{2.5} mass or its constituents. The authors note that these revised results are generally consistent with other long-term PM cohort studies, including the ACS and H6C studies.

¹ In addition to cohorts studies conducted in California, we have also identified a cross-sectional studies (Hankey et al., 2012). However, we have not summarized that study here.

Table 5.B-10. Summary of Effect Estimates From Associated With Change in Long-Term Exposure to PM_{2.5} in Recent Cohort Studies in California

Authors	Cohort	Hazard Ratios per 10 µg/m ³ Change in PM _{2.5} (95 th percentile confidence interval)		
		All Causes	Cardiopulmonary	Ischemic Heart Disease
McDonnell et al. (2000) ^a	Adventist Health Study (AHS) cohort (age > 27)	1.09 (.98–1.24)	N/A	N/A
Jerrett et al. (2005) ^b	Subset of the ACS cohort living in the Los Angeles metropolitan area (age > 30)	1.15 (1.03–1.29)	1.10 (0.94–1.28)	1.32 (1.03–1.29)
Chen et al. (2005) ^c	Adventist Health Study (AHS) cohort living in San Francisco, South Coast (i.e., Los Angeles and eastward), and San Diego air basins (age > 25)	N/A	N/A	1.42 (1.06–1.90)
Enstrom et al. (2005) ^d	California Prevention Study (age >65)	1.04 (1.01–1.07)	N/A	N/A
Krewski et al. (2009) ^e	Subset of the ACS cohort living in the 5-county Los Angeles Metropolitan Statistical Area (age > 30)	1.42 (1.26–1.27)	1.11 (0.95–1.23)	1.32 (1.06–1.64)
Ostro et al. (2010) ^c	California Teacher’s study. Current and former female public school professionals (age > 22)	1.84 (1.66–2.05)	2.05 (1.80–2.36)	2.89 (2.27–3.67)
Ostro et al. (2011) ^{c,f}		1.06 (0.96–1.16)	1.19 (1.05–1.36)	1.55 (1.24–1.93)

^a Table 3, adjusted for 10 µg/m³ change in PM_{2.5}.

^b Table 1. 44 individual-level co-variates + all social (i.e., ecologic) factors specified (principal component analysis).

^c Women only.

^d Represents deaths occurring from 1973–1982, but no significant associations were reported with deaths in later time periods. The PM ISA (U.S. EPA, 2009) concludes that the use of average values for California counties as exposure surrogates likely leads to significant exposure error, as many California counties are large and quite topographically variable.

^e Table 23. 44 individual-level co-variates + all social (i.e., ecologic) factors specified.

^f Erratum Table 2.

As shown in Table 5.B.10, most of the cohort studies conducted in California report central effect estimates similar to the (nation-wide) all-cause mortality risk estimate we applied from Krewski et al. (2009) and Laden et al. (2006) albeit with wider confidence intervals. A couple cohort studies conducted in California indicate higher risks than the risk estimates we applied.

5.B.6 References

- Abbey, David E et al. 1995. "Chronic Respiratory Symptoms Associated with Estimated Long-Term Ambient Concentrations of Fine Particulates less than 2.5 Microns in aerodynamic diameter (PM_{2.5}) and Other Air Pollutants." *Journal of Exposure Analysis and Environmental Epidemiology* 5(2): 137-158.
- Abt Associates, Inc. 2010. *BenMAP User's Manual Appendices*. Prepared for U.S. Environmental Protection Agency Office of Air Quality Planning and Standards. Research Triangle Park, NC. Available on the Internet at <<http://www.epa.gov/oaqps001/benmap/models/BenMAPManualAppendicesAugust2010.pdf>>.
- Abt Associates, Inc. 2011. *BenMAP User's Manual Appendices*. Prepared for U.S. Environmental Protection Agency Office of Air Quality Planning and Standards. Research Triangle Park, NC. September. Available on the Internet at <<http://www.epa.gov/air/benmap/models/BenMAPAppendicesSeptember2011.pdf>>.
- Agency for Healthcare Research and Quality (AHRQ). 2000. *HCUPnet, Healthcare Cost and Utilization Project*. Rockville, MD. Available on the Internet at <<http://hcupnet.ahrq.gov>>.
- Agency for Healthcare Research and Quality (AHRQ). 2007. *HCUPnet, Healthcare Cost and Utilization Project*. Rockville, MD. Available on the Internet at <<http://hcupnet.ahrq.gov>>.
- Chen LH, Knutsen SF, Shavlik D, Beeson WL, Petersen F, Ghamsary M, et al. 2005. "The association between fatal coronary heart disease and ambient particulate air pollution: are females at greater risk?" *Environ Health Perspect* 113:1723–1729.
- Clancy, L., P. Goodman, H. Sinclair, and D.W. Dockery. 2002. "Effect of Air-pollution Control on Death Rates in Dublin, Ireland: An Intervention Study." *Lancet* Oct 19;360(9341):1210-4.
- Desvousges, W.H., F.R. Johnson, and H.S. Banzhaf. 1998. *Environmental Policy Analysis With Limited Information: Principles and Applications of the Transfer Method* (New Horizons in Environmental Economics.) Edward Elgar Pub: London.

- Dockery, D.W., C.A. Pope, X.P. Xu, J.D. Spengler, J.H. Ware, M.E. Fay, B.G. Ferris, and F.E. Speizer. 1993. "An Association between Air Pollution and Mortality in Six U.S. Cities." *New England Journal of Medicine* 329(24):1753-1759.
- Enstrom JE 2005. "Fine particulate air pollution and total mortality among elderly Californians, 1973-2002." *Inhal Toxicol* 17: 803-816.
- Hankey, S., J. D. Marshall, et al. (2011). "Health Impacts of the Built Environment: Within-Urban Variability in Physical Inactivity, Air Pollution, and Ischemic Heart Disease Mortality." *Environ Health Perspect* 120(2).
- Jerrett M, Burnett RT, Ma R, Pope CA III, Krewski D, Newbold KB, et al. 2005. "Spatial analysis of air pollution and mortality in Los Angeles." *Epidemiology* 16(6):727-736.
- Kleckner, N., and J. Neumann. 1999. "Recommended Approach to Adjusting WTP Estimates to Reflect Changes in Real Income." Memorandum to Jim Democker, US EPA/OPAR. June 3, Available on the Internet at <<http://www.epa.gov/ttn/ecas/regdata/Benefits/background/klecknerandneumann1999.pdf>>.
- Krewski D, Jerrett M, Burnett RT, Ma R, Hughes E, Shi, Y, et al. 2009. "Extended follow-up and spatial analysis of the American Cancer Society study linking particulate air pollution and mortality." *HEI Research Report, 140*, Health Effects Institute, Boston, MA.
- Laden F, Schwartz J, Speizer FE, Dockery DW. 2006. "Reduction in fine particulate air pollution and mortality: extended follow-up of the Harvard Six Cities study." *Am J Respir Crit Care Med* 173:667-672.
- McDonnell WF; Nishino-Ishikawa N; Petersen FF; Chen LH; Abbey DE. 2000. "Relationships of mortality with the fine and coarse fractions of long-term ambient PM10 concentrations in nonsmokers." *J Expo Sci Environ Epidemiol*, 10: 427-436.
- Metzger, K. B., P. E. Tolbert, et al. (2004). "Ambient air pollution and cardiovascular emergency department visits." *Epidemiology* 15(1): 46-56.
- Miller, Kristin A., David S. Siscovick, Lianne Sheppard, Kristen Shepherd, Jeffrey H. Sullivan, Garnet L. Anderson, and Joel D. Kaufman. 2007. "Long-Term Exposure to Air Pollution and Incidence of Cardiovascular Events in Women." *New England Journal of Medicine*. 356 (5) :447-458.
- Ostro B, Lipsett M, Reynolds P, Goldberg D, Hertz A, Garcia C, et al. 2010. "Long-Term Exposure to Constituents of Fine Particulate Air Pollution and Mortality: Results from the California Teachers Study." *Environ Health Perspect* 118:363-369.
- Ostro B, Reynolds P, Goldberg D, Hertz A, et al. 2011. "Assessing Long-Term Exposure in the California Teachers Study." *Environ Health Perspect* June; 119(6): A242-A243.

- Peters, A., D. W. Dockery, J. E. Muller and M. A. Mittleman. 2001. "Increased particulate air pollution and the triggering of myocardial infarction." *Circulation*. Vol. 103 (23): 2810-5.
- Pope, C.A., III, M.J. Thun, M.M. Namboodiri, D.W. Dockery, J.S. Evans, F.E. Speizer, and C.W. Heath, Jr. 1995. "Particulate Air Pollution as a Predictor of Mortality in a Prospective Study of U.S. Adults." *American Journal of Respiratory Critical Care Medicine* 151:669-674.
- Röösli M, Kunzli N, Braun-Fahrlander C. 2004. "Use of Air Pollution 'Intervention-Type' Studies in Health Risk Assessment." *16th Conference of the International Society for Environmental Epidemiology*, New York, August 1-4, 2004.
- Röösli M, Kunzli N, Braun-Fahrlander C, Egger M. 2005. "Years of life lost attributable to air pollution in Switzerland: dynamic exposure-response model." *International Journal of Epidemiology* 34(5):1029-35.
- Tolbert, P. E., M. Klein, et al. 2007. "Multipollutant modeling issues in a study of ambient air quality and emergency department visits in Atlanta." *J Expo Sci Environ Epidemiol* 17 Suppl 2: S29-35.
- U.S. Environmental Protection Agency—Science Advisory Board (EPA-SAB). 1999a. *The Clean Air Act Amendments (CAAA) Section 812 Prospective Study of Costs and Benefits (1999): Advisory by the Health and Ecological Effects Subcommittee on Initial Assessments of Health and Ecological Effects. Part 1*. EPA-SAB-COUNCIL-ADV-99-012. July. Available on the Internet at [http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/F64A21824D19766885257193005E51CA/\\$File/conadv12.pdf](http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/F64A21824D19766885257193005E51CA/$File/conadv12.pdf).
- U.S. Environmental Protection Agency—Science Advisory Board (EPA-SAB). 1999b. *The Clean Air Act Amendments (CAAA) Section 812 Prospective Study of Costs and Benefits (1999): Advisory by the Health and Ecological Effects Subcommittee on Initial Assessments of Health and Ecological Effects. Part 2*. EPA-SAB-COUNCIL-ADV-00-001. October. Available on the Internet at [http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/7D54675178C12703852571B90044B4F8/\\$File/coua0001.pdf](http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/7D54675178C12703852571B90044B4F8/$File/coua0001.pdf).
- U.S. Environmental Protection Agency—Science Advisory Board (EPA-SAB). 2001. *Review of the Draft Analytical Plan for EPA's Second Prospective Analysis—Benefits and Costs of the Clean Air Act 1990-2020: An Advisory by a Special Panel of the Advisory Council on Clean Air Compliance Analysis*. EPA-SAB-COUNCIL-ADV-01-004. September. Available on the Internet at [http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/572DC1989A3986CC8525718D006BAB8B/\\$File/councila01004.pdf](http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/572DC1989A3986CC8525718D006BAB8B/$File/councila01004.pdf).

U.S. Environmental Protection Agency—Science Advisory Board (EPA-SAB). 2004. *Advisory on Plans for Health Effects Analysis in the Analytical Plan for EPA's Second Prospective Analysis—Benefits and Costs of the Clean Air Act, 1990-2020: Advisory by the Health Effects Subcommittee of the Advisory Council on Clean Air Compliance Analysis*. EPA-SAB-COUNCIL-ADV-04-002. March. Available on the Internet at <[http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/08E1155AD24F871C85256E5400433D5D/\\$File/council_adv_04002.pdf](http://yosemite.epa.gov/sab%5CSABPRODUCT.NSF/08E1155AD24F871C85256E5400433D5D/$File/council_adv_04002.pdf)>.

U.S. Environmental Protection Agency Science Advisory Board (U.S. EPA-SAB). 2010. *Review of EPA's DRAFT Health Benefits of the Second Section 812 Prospective Study of the Clean Air Act*. EPA-COUNCIL-10-001. June. Available on the Internet at <<http://yosemite.epa.gov/sab/sabproduct.nsf/9288428b8eeea4c885257242006935a3/59e06b6c5ca66597852575e7006c5d09!OpenDocument&TableRow=2.3#2.>>>.