

Health Practices and Cancer Mortality Among Active California Mormons

James E. Enstrom*

Religiously active Mormons in California are a nonsmoking population with unusually low risk for cancer. This finding is based on the results of our 1979 questionnaire survey of life-style and the 8-year (1980-1987) follow-up of mortality among 5,231 Mormon high priests and 4,613 wives 25-99 years of age. Our study, which is the first prospective cohort study of Mormons, shows low standardized mortality ratios (SMRs) for this population, relative to those for whites in the general population in the United States, which are defined as 100. The SMRs for males are 47 for all cancers, 52 for cardiovascular diseases, and 47 for all causes; the SMRs for females are 72 for all cancers, 64 for cardiovascular diseases, and 66 for all causes. For middle-aged high priests adhering to three health practices (never smoking cigarettes, engaging in regular physical activity, and getting proper sleep), the SMRs are 34 for all cancers, 14 for cardiovascular diseases, and 22 for all causes. These results have been largely replicated in an active Mormon-like subgroup (white nonsmokers attending church weekly) from a representative sample of residents of Alameda County, CA. Our findings confirm and expand on previous descriptive studies of Mormons and demonstrate how these results can be generalized. [J Natl Cancer Inst 81:1807-1814, 1989]

One means of gaining new insights into cancer etiology and prevention is to examine a well-defined low-risk population within the United States. It has been conclusively demonstrated (1-8) that members of the Church of Jesus Christ of Latter-day Saints, more popularly known as Mormons, constitute one such low-risk group. The Mormon church has about 4 million members in the United States and 6.5 million members worldwide, including about 500,000 in California and 1.2 million in Utah (9).

This population is interesting from an epidemiologic standpoint because its "Word of Wisdom" advises against the use of tobacco, alcohol, coffee, tea, and drugs (1). Also, the religion emphasizes a strong family life and education, advocates good health practices in general, and recommends a well-balanced diet (1).

Religiously active Mormons, specifically high priests, essentially abstain from tobacco, alcohol, and caffeine. We have studied this group because their cancer mortality rates are substantially lower than those of the general population and those of Mormons as a whole—a finding demonstrated in our previous studies (2-4) and confirmed by other investi-

gators (8). High priests are long-term devout Mormons who have risen to the highest rank of the church's lay priesthood, a rank they retain for the rest of their lives. Many of them currently serve on a voluntary basis as local church leaders. Their wives hold no priesthood level, but most are active Mormons as well. The number of high priests in California was about 16,000 as of January 1, 1980, and 25,000 as of January 1, 1988.

Previous research on cancer in Mormons has dealt with mortality in California and Utah (1-4) and incidence in Utah (6-8). These descriptive results have been based on church membership statistics, church death records, and records of newly diagnosed cases of cancer identified by religious affiliation. Life-style data on small samples of Mormons have confirmed that active Mormons do not smoke. However, abstinence from smoking does not fully explain the descriptive cancer mortality patterns observed (2). Consequently, we undertook a prospective epidemiologic study of a well-defined cohort of active Mormons.

This study provides the first detailed examination and explanation of the life-style characteristics of Mormons in relation to subsequent mortality rates. This detailed long-term assessment of the life-style of active Mormons as it relates to cancer mortality and minimum cancer risk provides new information regarding the actual prevention of cancer in a population in the United States.

Methods

Questionnaire

A four-page questionnaire to measure the life-style characteristics of active Mormons was prepared at the University of California, Los Angeles (UCLA). This questionnaire was

Received May 1, 1989; revised August 18, 1989; accepted August 31, 1989.

Supported by American Cancer Society grants PDT-332 and PDT-51AB; and by Public Health Service grant CA-00748 (Preventive Oncology Academic Award) from the National Cancer Institute, National Institutes of Health, Department of Health and Human Services.

School of Public Health and Jonsson Comprehensive Cancer Center, University of California, Los Angeles, CA.

I thank Lester Breslow and the Human Population Laboratory for access to the Alameda County data, and Linda E. Kanim, Richard J. Biermann, and Michael Succar for technical assistance.

*Correspondence to: Dr. James E. Enstrom, School of Public Health, University of California, 10833 Le Conte Ave., Los Angeles, CA 90024.

sent around December 1, 1979, by Mormon church personnel, to high priests and their wives in California. To maintain the confidentiality of the members unwilling to participate, our questionnaires were mailed from the church headquarters in Salt Lake City, UT, and willing respondents replied directly to UCLA. A letter accompanying the questionnaire made clear that the study was being done independently of the church.

A one-time mailing of two questionnaires (one for husband and one for wife) was made to about 12,000 California households of high priests with valid addresses on the church's computerized membership file as of about December 1, 1979. Responses from 9,844 persons 25-99 years of age (5,231 high priests and 4,613 wives) were received at UCLA. The church did not permit additional mailings to nonrespondents.

The completed questionnaires provided data on basic demographic characteristics; major components of life-style, diet, and medical history; and other health-related characteristics. It incorporated parts of a 1974 Health and Ways of Living questionnaire used in Alameda County by the Human Population Laboratory of the California State Department of Health Services (10).

Mortality Follow-up

For all California high priests, including those in our cohort, complete and accurate mortality data have been obtained for 8 years, from January 1, 1980, through December 31, 1987. We have obtained 1980-1987 mortality data on all high priests in California from the Mormon church computerized membership file in Salt Lake City. To verify deaths and determine the underlying cause of death, we computer matched the church data on the deceased high priests with the data on all California deaths, matching on the full name, date and place of birth, and date and place of death of the deceased person. The underlying cause for each California death has been assigned by the state nosologist, following the rules of the ninth revision of the *International Classification of Diseases* (ICD9). The mortality data and the age-specific membership statistics obtained from the Mormon church have been used to calculate mortality rates for all Mormon high priests in California during 1980-1987. These mortality rates update previously published rates for 1968-1975 (3). Similarly, the data from the high priest cohort have been computer matched with the 1980-1987 mortality data obtained from the files of the Mormon church and the state of California to identify all cohort deaths. We estimate that no more than 2% of all deaths in cohort members in California have been missed by matching with these two sources. We have identified 612 deaths of cohort members in California, but deaths of members who left the state have not been included.

We have used the California Department of Motor Vehicles (DMV) files to verify the vital status and current addresses of the high priests and to estimate their movement out of California. In 1980, 98% of cohort members and 99% of households were linked with the DMV driver's license file. This file has been updated twice a year with notation made when individuals left California for another state or died. As

of January 1, 1988, approximately 11% of the cohort members had left California, according to the DMV.

Comparison Group

To compare the high priest cohort with the general population, we made use of data on the 1974 representative sample of 3,119 noninstitutionalized adults in Alameda County, which has been used previously to study the relationship between health practices and mortality (10) and to make comparisons with Mormons (4). Alameda County is similar to the state of California and the United States as a whole in terms of population demographics and mortality rates.

Analysis

Our analysis of life-style is based on three general health practices:

- (1) never smoked cigarettes;
- (2) engage in regular physical activity (often or sometimes engage in active sports, swim, or take long walks or often garden or do physical exercises); and
- (3) get proper sleep (usually 7-8 hr/day).

Each of these health practices has repeatedly been related to total mortality rates, in other studies (10-14).

Standardized mortality ratios (SMRs) for major cancer sites, all cancers (ICD9 140-208), all cardiovascular diseases (ICD9 390-459), and all causes are calculated for the total cohort as well as for the health practice subgroups. SMRs are presented for persons in the 25-99-year age group during the study period and for persons in the 24-64-year age group at entry in the study.

The SMR for each subgroup is the number of observed deaths divided by the number of expected deaths, expressed as a percentage. The total number of expected deaths is the sum of the expected deaths for each 5-year age group. The number of expected deaths for one 5-year age group is the number of person-years of risk multiplied by the age-specific mortality rate for that age group of whites in the United States. The total person-years at risk are based on the person-years at risk among California residents. For each person, the period of risk began January 1, 1980, and ended on the date of death; date of leaving California according to the DMV; or December 31, 1987, whichever came first.

We calculated each SMR and its 95% confidence interval using a survival program developed by Monson (15), with expected deaths based on concurrent death rates among whites in the United States. By this definition, the SMRs for whites in the general population in the United States are 100. We chose this population as the standard for comparison of mortality because detailed data are widely available and the mortality rates are similar to those for comparable groups, such as California whites or Alameda County whites. Because of the small number of deaths in some subgroups, SMRs, which are based on indirect age standardization, have been used instead of age-adjusted death rates, which are based on direct age standardization. However, the results by both methods are consistent. The age distribution in the high priest cohort at the time of enrollment was as follows: 7% for ages 25-34 years, 21% for ages 35-44, 28% for ages 45-54, 23% for

ages 55-64, 14% for ages 65-74, 6% for ages 75-84, and 1% for ages 85-99.

Results

Several basic demographic and health-related characteristics of the 1979 California high priest cohort and the 1974 Alameda County representative cohort are shown in table 1 for males and in table 2 for females. These persons were 25-64 years of age at the time of entry. Data are presented for all whites in the Alameda County cohort and for two white subgroups: Mormons who attend church weekly and non-Mormon nonsmokers (former smokers and those who never smoked) who attend church weekly. The Alameda County Mormons were identified on the basis of their answers to the questions on religion and church attendance. There are only 27 of them, but within statistical fluctuation, their characteristics are similar to those of the high priest

cohort. The high priests are white, married, and well educated; attend church weekly; and use essentially no tobacco, alcohol, or caffeine.

Table 3 shows that the SMRs during 1980-1987 for an average of 21,000 California high priests are 50 for all cancers, 24 for smoking-related cancers, 69 for non-smoking-related cancers, and 53 for all causes. The SMRs during 1980-1987 for 5,231 high priests from the 1979 cohort are 47 for all cancers, 15 for smoking-related cancers, 71 for non-smoking-related cancers, and 47 for all causes. The results for the high priest cohort are similar to those for all California high priests. The difference for lung cancer (7 vs. 20) may be due to the fact that there is a somewhat larger proportion of persons who never smoked in the cohort than among all California high priests. The small difference for all causes (47 vs. 53) may be due to a small questionnaire-respondent bias (see Discussion section).

Table 4 shows the SMRs for all causes, all cancers, and

Table 1. Major demographic and health-related characteristics of males in 1979 California Mormon high priest cohort and 1974 Alameda County general population cohort 25-64 yr of age*

Characteristic	1979 California Mormon high priest cohort	1974 Alameda County whites		
		Active Mormons†	Religious non-Mormon nonsmokers‡	Total
Sample size	4,112	11	91	788
Race (white)	99	100	100	100
Age (mean in yr)	48.0	37.1	45.6	41.8
Church attendance weekly	99	100	100	17
"Church group" members	100	100	71	20
Marital status				
Married	99	82	92	82
Never married	0	18	3	10
Education (mean in yr)	16.1	15.3	12.9	13.6
<12 yr	3	9	21	16
≥16 yr	63	64	23	32
Occupation (professional, technical, or managerial)	73	73	41	48
Height (mean in inches)	70.5	70.7	70.4	70.4
Weight (mean in lb)	184	184	180	177
Use of tobacco, alcohol, or caffeine				
Cigarettes				
Current	0	9	0	39
Former	25	36	46	29
Never	75	55	54	32
Cigars (current)	0	0	5	7
Pipe (current)	0	0	3	11
Beer (current)	0	0	67	75
Wine (current)	1	0	69	77
Liquor (current)	0	0	65	79
Coffee (current)	1	0	82	83
Tea (current)	1	0	24	20
Active sports				
Often	19	18	11	18
Sometimes	43	64	59	52
Physical exercises				
Often	29	9	22	21
Sometimes	47	64	52	49
Sleep, 7-8 hr	82	73	82	79
Current health status				
Excellent	55	55	44	38
Good	40	45	51	53
Fair	4	0	4	7
Poor	1	0	1	2

* Unless specified otherwise, values = %.

† Mormons who attend church weekly.

‡ Non-Mormon nonsmokers who attend church weekly.

Table 2. Major demographic and health-related characteristics of females of 1979 California Mormon high priest cohort and 1974 Alameda County general population cohort 25-64 yr of age*

Characteristic	1979 California Mormon high priest cohort	1974 Alameda County whites		
		Active Mormons†	Religious non-Mormon nonsmokers‡	Total
Sample size	3,968	16	148	878
Race (white)	100	100	100	100
Age (mean in yr)	46.4	35.3	43.6	42.5
Church attendance weekly	99	100	100	23
"Church group" members	100	100	66	24
Marital status				
Married	100	88	81	76
Never married	0	12	5	6
Education (mean in yr)	13.8	13.1	12.6	13.1
<12 yr	4	6	33	16
≥16 yr	26	25	18	23
Occupation (professional, technical, or managerial)	10	19	27	31
Height (mean in inches)	64.6	65.3	64.4	64.5
Weight (mean in lb)	146	139	138	138
Use of tobacco, alcohol, or caffeine				
Cigarettes				
Current	0	0	0	37
Former	9	19	24	17
Never	91	81	76	46
Cigars (current)	0	0	0	0
Pipe (current)	0	0	0	0
Beer (current)	0	0	26	41
Wine (current)	0	0	60	70
Liquor (current)	0	0	55	70
Coffee (current)	1	6	74	79
Tea (current)	2	6	46	37
Active sports				
Often	5	6	6	10
Sometimes	28	44	39	41
Physical exercises				
Often	23	19	24	20
Sometimes	52	56	51	52
Sleep, 7-8 hr	84	81	84	78
Current health status				
Excellent	45	31	41	35
Good	46	56	49	51
Fair	7	13	8	12
Poor	1	0	2	2

*Unless specified otherwise, values = %.

†Mormons who attend church weekly.

‡Non-Mormon nonsmokers who attend church weekly.

Table 3. SMRs for major cancer sites and all causes of cancer in California Mormon high priests and their wives 25-99 yr of age*

Cause of death (ICD9 classification)	All California high priests (males, n = 21,000)		1979 California high priest cohort (males, n = 5,231)		1979 California high priest cohort (females, n = 4,613)	
	Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)
Colorectal cancer (153-154)	68	61 (48-78)	16	65 (37-105)	10	88 (42-162)
Lung cancer (162)	64	20 (16-26)	5	7 (2-16)	8	43 (18-85)
Breast cancer (174)	—	—	—	—	18	90 (54-143)
Prostate cancer (185)	95	100 (81-123)	22	104 (65-157)	—	—
Smoking-related cancers (140-150, 162, 188)	96	24 (20-30)	13	15 (8-26)	9	41 (19-78)
Non-smoking-related cancers	375	69 (62-76)	85	71 (57-88)	60	82 (63-107)
All cancers (140-208)	471	50 (46-55)	98	47 (38-57)	69	72 (56-92)
All cardiovascular diseases (390-459)	1,157	58 (55-61)	230	52 (45-59)	89	64 (52-79)
All causes	2,091	53 (51-55)	406	47 (42-52)	206	66 (58-76)

*Based on church statistics for an average of 21,000 California high priests during 1980-1987 and for 5,231 California high priests and 4,613 wives in the 1979 cohort who had follow-up during 1980-1987. 95% confidence interval (CI) is based on a Poisson-distributed variable.

Table 4. Results at 8-yr follow-up (1980-1987) of 1979 cohort of California Mormon high priests and their wives 25-99 yr of age: SMRs for major causes of death, by sex and health practices*

Health practices	No. alive Jan. 1, 1980	All causes		All cancers		All cardiovascular diseases	
		Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)
Males							
Total	5,231	406	47 (42-52)	98	47 (38-57)	230	52 (45-59)
1 HP	3,643	206	40 (35-46)	58	46 (35-60)	116	45 (37-54)
2 HP	3,419	164	35 (30-41)	53	46 (34-60)	87	37 (30-46)
3 HP	2,802	124	33 (28-40)	45	48 (35-65)	62	34 (26-43)
Females							
Total	4,613	206	66 (58-76)	69	72 (56-92)	89	64 (52-79)
1 HP	4,152	188	68 (59-79)	60	71 (54-91)	81	65 (52-81)
2 HP	3,525	130	58 (49-69)	45	64 (47-86)	54	55 (41-72)
3 HP	2,942	90	52 (41-63)	32	56 (39-80)	34	45 (31-64)
Both sexes							
Total	9,844	612	52 (48-56)	167	55 (47-64)	319	55 (49-66)
1 HP	7,795	394	50 (45-55)	118	56 (46-67)	197	52 (45-60)
2 HP	6,944	294	43 (38-48)	98	53 (43-66)	141	43 (35-51)
3 HP	5,744	214	39 (34-45)	77	51 (40-64)	96	37 (30-46)

*Health practices are as follows: 1 HP = never smoked cigarettes; 2 HP = never smoked cigarettes and engage in regular physical activity; 3 HP = never smoked cigarettes, engage in regular physical activity, and get proper sleep (7-8 hr/day). Regular physical activity = often or sometimes engage in active sports, swim, or take long walks or often garden or do physical exercises.

all cardiovascular diseases for persons 25-99 years of age in the high priest cohort, by sex and the three health practices. The SMRs generally declined with adherence to more of these practices. The SMRs for all causes declined from 47 to 33 for males and from 66 to 52 for females. The SMRs for all cancers changed from 47 to 48 for males and from 72 to 56 for females. The SMRs for all cardiovascular diseases declined from 52 to 34 for males and from 64 to 45 for females. Several health practices other than the three we considered in our analyses were related to mortality in the Alameda County study (10,11), but they had no additional impact on the SMRs for the high priest cohort. These additional practices were maintenance of proper weight for

height, moderate alcohol consumption, eating breakfast, and not eating between meals. We have not yet explored all possible health and dietary practices in relationship to mortality, so it is possible that other combinations could have an impact on the SMRs.

Table 5 shows the SMRs for persons in the high priest cohort who were 25-64 years of age at entry in the study. These SMRs are lower than those shown in table 4 for persons in the 25-99-year age group. The differences are large for males and small for females. For ages 25-64 years, the SMRs for males are 30 for all cancers and 28 for all causes, and the SMRs for females are 76 for all cancers and 62 for all causes. For persons adhering to the three health

Table 5. Results at 8-yr follow-up (1980-1987) of 1979 cohort of California Mormon high priests and their wives 25-64 yr of age at entry in study: SMRs for major causes of death, by sex and health practices*

Health practices	No. alive Jan. 1, 1980	All causes		All cancers		All cardiovascular diseases	
		Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)
Males							
Total	4,112	95	28 (22-34)	30	30 (21-43)	41	26 (19-36)
1 HP	3,060	56	23 (18-30)	22	32 (20-49)	18	17 (10-26)
2 HP	2,913	53	23 (17-31)	22	34 (21-52)	17	17 (10-27)
3 HP	2,403	40	22 (15-29)	18	34 (20-54)	12	14 (7-25)
Females							
Total	3,968	95	62 (50-76)	48	76 (57-102)	22	43 (27-66)
1 HP	3,589	84	61 (49-76)	41	73 (53-100)	19	42 (25-66)
2 HP	3,093	63	53 (41-68)	31	65 (44-92)	16	41 (23-67)
3 HP	2,610	46	47 (34-62)	22	55 (35-84)	11	34 (17-61)
Both sexes							
Total	8,080	190	38 (33-44)	78	48 (38-60)	63	30 (23-39)
1 HP	6,649	140	37 (31-44)	63	51 (40-66)	37	24 (17-33)
2 HP	6,006	116	34 (28-41)	53	47 (35-62)	33	23 (16-32)
3 HP	5,013	86	30 (24-37)	40	43 (31-58)	23	20 (13-30)

*For definitions, see footnote to table 3.

practices, the SMRs for all causes declined even further: from 28 to 22 for males and from 62 to 47 for females. The SMRs for all cancers did not vary significantly with the number of health practices, but the SMRs for all cardiovascular diseases declined substantially: from 26 to 14 for males and from 43 to 34 for females.

To see the extent to which these results can be reproduced in the general population, we analyzed the 1974 representative sample of noninstitutionalized Alameda County residents (10) for residents with characteristics similar to those of active Mormons. We defined "active Mormon-like" persons as residents of all religions who are white, do not smoke cigarettes (former smokers and those who never smoked), and attend church weekly. Their SMRs for all causes were 42 for males and 61 for females, as shown in table 6. The SMRs for all whites, whites who never smoked cigarettes, active Mormon-like persons, and active Mormon-like persons with the three health practices are presented in table 6 for all causes, all cancers, and all cardiovascular diseases. The SMRs for active Mormon-like persons in the Alameda County cohort are the same as the corresponding SMRs for the high priest cohort (table 4), within statistical fluctuation. Although the data are not shown in table 6, the SMRs for all Alameda County whites were substantially less than 100 during the first 5 years of follow-up, but thereafter, the values were about 100, as the effect of initially including only noninstitutionalized persons diminished.

Logistic regression analyses of data from the Alameda County cohort show that the associations of mortality with age, sex, initial health status, cigarette smoking status, and frequency of church attendance are statistically significant ($P < .05$). Several additional variables, such as race, mar-

ital status, education level, and other health practices were not significant in the regression analyses after the first five variables were accounted for. Thus, the overall SMRs for the high priest cohort appear to be reasonably well explained by the fact that the cohort consists of white nonsmokers who attend church weekly. The analyses are not exhaustive, however, and other variables may be significant. For instance, weekly church attendance may be related to other variables yet to be determined.

Discussion

These findings indicate that active Mormons (high priests and their wives) have very low rates of mortality from all cancers and all causes. The plausibility of these findings is supported by related research. Church attendance has previously been associated with reduced mortality in a sample of the general population in Maryland (16,17) and in a few other populations (18). For many decades, reduced mortality rates have been reported among clergy (19,20) and among health-conscious religious groups, such as Seventh-Day Adventists (21) and Mormons (1). Although high priests are lay clergymen, they may be similar in many ways to formal clergymen. Also, these results may be partially influenced by the strong social network ties among high priests, who are all married and belong to a church group. Social networks have previously been related to total mortality, independently of other health practices (22,23).

Table 7 shows a comparison of SMRs for all cancers and all causes in major low-risk populations in the United States. These populations are whites who never smoked, healthy questionnaire respondents who never smoked, nonsmoking

Table 6. Results at 14-yr follow-up (1974-1987) of 1974 representative sample of noninstitutionalized Alameda County population: SMRs for major causes of death, by sex, smoking status, active Mormon-like characteristics, and health practices*

Subgroup	No. alive Jan. 1, 1974	All causes		All cancers		All cardiovascular diseases	
		Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)	Observed deaths	SMR (95% CI)
Males							
All whites	1,036	151	81 (68-94)	34	80 (55-111)	76	79 (62-98)
Whites who never smoked cigarettes	364	50	69 (51-91)	9	58 (26-110)	29	76 (51-110)
Active Mormon-like persons	136	14	42 (23-71)	4	51 (14-131)	7	41 (16-84)
Active Mormon-like persons with 3 HP	57	3	22 (4-65)	0	0 (0-116)	3	44 (9-127)
Females							
All whites	1,254	159	88 (75-103)	39	99 (70-135)	77	77 (61-97)
Whites who never smoked cigarettes	627	84	70 (56-87)	13	59 (31-100)	42	59 (42-79)
Active Mormon-like persons	233	28	61 (40-88)	5	54 (17-125)	15	56 (31-93)
Active Mormon-like persons with 3 HP	130	10	48 (23-88)	1	21 (0-117)	6	53 (19-116)
Both sexes							
All whites	2,290	310	84 (75-94)	73	89 (70-112)	153	78 (66-92)
Whites who never smoked cigarettes	991	134	70 (59-83)	22	58 (36-88)	71	65 (51-83)
Active Mormon-like persons	369	42	53 (38-70)	9	52 (24-99)	22	50 (31-76)
Active Mormon-like persons with 3 HP	187	13	38 (20-65)	1	13 (1-70)	9	49 (22-93)

* Active Mormon-like persons are white nonsmokers of all religions who attend church weekly. These include active Mormons and religious non-Mormon nonsmokers in tables 1 and 2. HP = health practices.

Table 7. Preventability of cancer on basis of SMRs for all cancers and all causes of death among various low-risk white populations in United States*

Comparison group	SMR			
	All cancers		All causes	
	Males	Females	Males	Females
Ages 25-99 yr				
U.S. whites	100	100	100	100
U.S. whites who never smoked†	65	86	74	84
Healthy questionnaire respondents who never smoked†	~50	~82	~55	~75
Nonsmoking members of health-conscious religious groups†	~50	~70	~50	~70
Mormon high priest questionnaire respondents	47	72	47	66
Mormon high priest questionnaire respondents with 3 HP	48	56	33	52
Ages 25-64 yr				
U.S. whites	100	100	100	100
U.S. whites who never smoked†	57	78	65	75
Healthy questionnaire respondents who never smoked†	~45	~70	~50	~70
Nonsmoking members of health-conscious religious groups†	~35	~65	~35	~65
Mormon high priest questionnaire respondents	30	76	28	62
Mormon high priest questionnaire respondents with 3 HP	34	55	22	47

*SMRs are based on comparison with whites in general U.S. population. HP = health practices.

†See ref. 5.

members of health-conscious religious groups, Mormon high priest questionnaire respondents, and a subgroup of these high priests with the three health practices. Except for our data for high priests from this study, the data on these low-risk populations are derived from other major epidemiologic studies during the past 35 years (5). The SMRs are based on comparison of each low-risk population with whites in the United States. The SMRs for all high priests are in agreement with those for nonsmoking members of health-conscious religious groups. However, for high priests with the three health practices, the SMRs for cardiovascular diseases and all causes are among the lowest ever reported.

To strengthen the validity of our findings, we have carefully examined three factors that could cause spurious results: underascertainment of deaths, selection bias due to the number of healthy questionnaire respondents, and unique results applicable only to the cohort. First, there is virtually complete ascertainment of our data on California deaths, because we have obtained consistent and overlapping reporting of deaths from three independent sources: California death files, Mormon church death files, and California DMV death records. In addition, dates of birth given on the questionnaires are in agreement with those on the DMV records, indicating that the age distribution is correct. Also, movement of cohort members from California has been accounted for by careful monitoring of driver's license records.

Second, there is little evidence that the SMRs were unusu-

ally low because of bias due to the number of healthy questionnaire respondents. This selection bias occurs when only the healthiest persons respond to a questionnaire mailed to a target population. The result is extremely low SMRs in the first few years of follow-up, followed by SMRs that steadily increase until the bias is no longer significant. But for our cohort, the SMRs have remained fairly constant in years 1-8 of follow-up. The SMRs for all causes, by year, are 44, 48, 40, 42, 48, 49, 47, and 56 for males and 78, 63, 57, 42, 68, 61, 74, and 90 for females. In addition, the SMRs for all causes are stable over the 8-year follow-up period when analyzed by the three health practices and by age at entry in the study. Also, the SMRs for the prospectively followed high priest cohort are in good agreement with concurrent SMRs calculated independently from Mormon church death records and membership statistics for all high priests. In addition, the SMRs agree with previously published descriptive mortality data on high priests (2-4,8). We believe the respondent bias is small, because high priests as a whole are such a healthy group and because our cohort includes a representative sample of them, even though the response rate to the one questionnaire mailing was not particularly high.

Third, the findings in our high priest cohort have been replicated to a large extent in a representative sample of the general population. Thus, these findings are not unique to our cohort; they are generalizable.

Conclusions

For high priests, especially those who are middle aged, the SMRs for all cancers, for cardiovascular diseases, and for all causes are substantially below those for typical nonsmokers. Indeed, this population is currently achieving the 50% reduction in cancer mortality that the National Cancer Institute has set as a goal for the year 2000 (24). These results demonstrate substantial progress in prevention of cancer and other diseases in one well-defined population, and they suggest a life-style that could result in a major reduction in cancer mortality, as well as mortality in general. Further research might reveal that additional health practices result in an even greater reduction in mortality.

References

- (1) ENSTROM JE: Cancer mortality among Mormons. *Cancer* 36:825-841, 1975
- (2) ENSTROM JE: Cancer and total mortality among active Mormons. *Cancer* 42:1943-1951, 1978
- (3) ENSTROM JE: Cancer mortality among Mormons in California during 1968-75. *JNCI* 65:1073-1082, 1980
- (4) ENSTROM JE: Health and dietary practices and cancer mortality among California Mormons. In *Banbury Report 4: Cancer Incidence in Defined Populations* (Cairns J, Lyon JL, Skolnick M, eds). Cold Spring Harbor, NY: Cold Spring Harbor Laboratory, 1980, pp 69-90
- (5) ENSTROM JE, KANIM LE: Populations at low risk. In *Cancer Prevention in Clinical Medicine* (Newett GR, ed). New York: Raven Press, 1983, pp 49-78
- (6) LYON JL, KLAUBER MR, GARDNER JW, ET AL: Cancer incidence in Mormons and non-Mormons in Utah, 1966-1970. *N Engl J Med* 294:129-133, 1976
- (7) LYON JL, GARDNER JW, WEST DW: Cancer incidence in Mormons and non-Mormons in Utah during 1967-75. *JNCI* 65:1055-1061, 1980
- (8) GARDNER JW, LYON JL: Cancer in Utah Mormon men by lay priesthood level. *Am J Epidemiol* 116:243-257, 1982

